

Anemia Status of Adults in Dakkilavaripalem – A Rural Area of Nellore

¹Katari Kantha* & ²Arumugam Indira

¹Department of Community Health Nursing, Narayana College of Nursing, Nellore,
Andhra Pradesh, 524003, India.

²Principal, Narayana College of Nursing, Nellore, Andhra Pradesh, 524003, India.

ABSTRACT

Background: Anemia is a major public health problem in India. Although nearly three quarters of the Indian population live in rural areas, the epidemiology of anemia in rural settings is not well known. The purpose of this study was to determine the prevalence of anemia among adults in selected rural areas of Nellore.

Objective: The objective was to assess the prevalence of anemia among rural adults.

Methods: The participants in this cross-sectional study came from Dakkilavaripalem rural area of Nellore. A total of 115 adults were randomly selected. Trained investigators administered a standard questionnaire to each participant during a face-to-face interview and carried out Haemoglobin testing by the haemometer.

Results: The results show that, out of 115 adults with regard to the category of the Anemia 70 (60.9%) had Normal (>11 gm/dl) Haemoglobin, 10 (8.7%) had Mild Anemia (9- 11 gm/dl), 2 (1.7%) had Moderate Anemia (7- 9 gm/dl) and haemoglobin testing was not done to 33(28.7%) due to unavailability of subjects on three repeated visits.

Keywords: Adults, Anemia, Prevalence, Nellore.

1. Introduction

According to the World Health Organization (WHO), there are two billion people with anemia in the world and half of the anemia is due to iron deficiency¹. Anemia is a late indicator of iron deficiency, so it is estimated that the prevalence of iron deficiency is 2.5 times that of anemia. Anemia and iron deficiency lead to substantial physical productivity losses in adults². Iron deficiency during pregnancy is associated with maternal mortality, preterm labor, low birth-weight, and infant mortality². In children, iron deficiency affects cognitive and motor development and increases susceptibility to infections³.

Anemia is a major health problem in India. In the 2005- 2006 National Family Health Survey (NFHS-3), a household survey aimed at having national and state representative data on population health and nutrition; the prevalence of anemia was 70% in children aged 6–59 months, 55% in females aged 15–49 years, and 24% in males aged 15–49 years⁴. Although the NFHS-3 showed that the prevalence of anemia was higher in rural areas, there is a paucity of data about the epidemiology of anemia in rural settings⁵. The aim of this study is to describe the prevalence of anemia among adults in Dakkilavaripalem, a rural area of Nellore in Andhra Pradesh, India.

Surveillance of anemia is challenging, requiring simultaneous understanding of the epidemiology of its underlying causes. Focusing only on total anemia invites misinterpretation of trends because changes in severity may be missed. Estimates in high-risk populations suggest that total anemia prevalence may be as high as 50% to 80%, with as many as 10% to 20% having moderate to severe anemia. Prevalence is consistently higher in people with low socioeconomic status, low body weight, and in females who have recently given birth.

The research of Dr. Indira. A et al. ⁶ and other studies^{7,8, &9} shows the nutritional status of various categories of people of Nellore.

The aim of this study was to assess the prevalence of anemia among rural adults.

2. Detailed Research Plan

2.1 Research Approach: Quantitative Approach.

2.2 Research Design: Descriptive design.

2.3 Research Setting: The study was conducted at Dakkilavaripalem rural area of Nellore.

2.4 Sampling Technique: Convenience sampling technique.

2.5 Sample Size: A total of 115 samples were included in this study.

3. RESULTS AND DISCUSSION:

SL NO	DEMOGRAPHIC VARIABLE	FREQUENCY	PERCENTAGE
1	AGE a. 20-30 years b. 31-40 years c. 41-50 years d. 51-60 years	37 36 23 19	32.2% 31.3% 20.0% 16.5%
2.	GENDER a. Male b. Female	47 68	40.9% 51.1%
3.	EDUCATION a. Illiterate b. Inter/SSLC c. Under Graduate d. Post Graduate	71 36 3 5	61.7% 31.3% 2.6% 4.3%
4.	FAMILY a. Extended Family b. Joint Family c. Nuclear Family	2 36 77	1.7% 31.3% 66.9%
5	OCCUPATION a. Farmer b. Coolie c. Managerial job d. Teaching job e. Others	23 69 0 4 19	20.0% 60.0% 0.0% 3.5% 16.5%
6	INCOME a. Rs. 2000 – 4000/- b. Rs. 4001 – 6000/- c. Rs. 6001 – 8000/- d. None	79 17 9 9	68.7% 14.8% 7.8% 7.8%
7	FOOD PATTERN a. Vegetarian b. Non – vegetarian c. Mixed	1 63 51	0.9% 54.8% 44.3%

The study included 82 determinations of haemoglobin from 115 adults, of which 47(40.9%) were male and 68 (51.1%) were female. Out of 115 adults 37 (32.2%) belongs to 20-30 years, 36 (31.3%) belongs to 31-40 years, 23 (20.0%) belongs to 41-50 years, 19 (16.5%) belongs to 51-60 yrs. Regarding food pattern, Out of 115 adults 1 (0.9%) was vegetarian, 63(54.8%) were non-vegetarian and 51(44.3%) were having mixed dietary practices.

The previous studies show that Spirulina was thought to be the best food for correcting the anemia¹⁰.

Table 2: Frequency and Percentage Distribution of Prevalence of Anemia in DAKKILIVARIPALEAM N=115

Classification	frequency	Percentage
Normal(>11 gm/dl)	70	60.9%
Mild(9- 11 gm/dl)	10	8.7%
Moderate(7- 9 gm/dl)	2	1.7%
Not done	33	28.7%




Fig 1: Percentage Distribution of Prevalence of Aneamia in DAKKILIVARIPALEAM

The above figure illustrates the anemia distribution in Dakkilavaripalem. Out of 115 adults with regard to the category of the Anemia 70 (60.9%) had Normal(>11 gm/dl) Haemoglobin, 10 (8.7%) had Mild Anemia (9- 11 gm/dl), 2 (1.7%) had Moderate Anemia (7- 9 gm/dl) and haemoglobin testing was not done to 33(28.7%) due to unavailability of subjects on three repeated visits.

In this study, we observed that majority 70 (60.9%) of the adults are having normal haemoglobin. Necessary actions need to be taken to correct the anemia for 12 (10.4%).

The present study results are consistent with the findings of the previous studies conducted in Nellore region^{11,12,13,14&15}.

Conclusion: The results of this study can be used by public health programme to design target interventions aimed at reducing the burden of anemia in Dakkilavaripalem rural area of Nellore. Further studies are needed to clarify the etiology of anemia among adults. This in turn might shed light on the preventive and intervention strategies that might be effective in this community.

References:

- [1] WHO, UNICEF, and UNU, Iron Deficiency Anaemia: Assessment, Prevention and Control, A Guide for Programme Managers, WHO, UNICEF, UNU, Geneva, Switzerland, 2001, http://www.who.int/nutrition/publications/micronutrients/anaemia_iron_deficiency/WHO_NHD_01.3/en/index.html.

- [2] M. B. Zimmermann and R. F. Hurrell, "Nutritional iron deficiency," *The Lancet*, vol. 370, pp. 511–520, 2007.
- [3] R. D. Baker, F. R. Greer, and Committee on Nutrition American Academy of Pediatrics, "Diagnosis and prevention of iron deficiency and iron-deficiency anemia in infants and young children (0–3 years of age)," *Pediatrics*, vol. 126, pp. 1040–1050, 2010.
- [4] F. Arnold, S. Parasuraman, P. Arokiasamy, and M. Kothari, "Nutrition in India," in *National Family Health Survey (NFHS-3) India 2005-06, 2009*, http://www.rchiips.org/nfhs/nutrition_report_for_website_18sep09.pdf.
- [5] Office of The Registrar General & Census Commissioner, Census of India, 2011.
- Arumugam Indira, Katari Kantha. Nutritional status of antenatal and postnatal mothers in selected rural areas at Nellore. *International Journal of Recent Scientific Research*. 2015; 6(2):2796-2800. **View Article Google Scholar**
- Katari Kantha *et al.*, Nutritional Status of infant Children in selected rural areas at Nellore international journal of recent scientific research. 2015; 6(6):4597- 4601. **View Article Google Scholar**
- Katari Kantha *et al.*, Knowledge of the Expectant Fathers Regarding safe Motherhood in Narayana Medical College Hospital, Nellore. *International Journal of Recent Scientific Research*. 2015; 6(4):3423- 3427. **View Article Google Scholar**
- Katari Kantha, Usha Rani M, Alfia Parameswaran, Arumugam Indira. The

knowledge regarding eating disorders among adolescent girls International Journal of Applied Research. 2016; 2(5):864-866. **View Article Google Scholar**

10. Kantha Katari. Spirulina the best food for future. www.scopemed.org/mno=228758 [Access: September 26, 2016]. **View Article Google Scholar**

11. Kantha Katari. Human milk banking. www.scopemed.org/mno=229095 [Access: September 26, 2016]. **View Article Google Scholar**

12. Kantha Katari. Knowledge regarding complementary and alternative therapy among staff nurses. www.scopemed.org/mno=229182 [Access: September 26, 2016]. **View Article Google Scholar**

13. Arumugam Indira and Katari Kantha. 2017, Correlation of Body Mass Index Levels With Age at Menarche in Women In, Selected Rural Areas of Nellore: A Cross Sectional Study. *Int J Recent Sci Res.* 8(4), pp. 16297-16299. DOI: http://dx.doi.org/10.24327/ijrsr.2017.0804_0122. **View Article Google Scholar**

14. E Vanitha Sree, Katari Kantha and Arumugam Indira Knowledge regarding gastrostomy feeding among staff nurses and nursing students International Journal of Applied Research 2017; 3(3): 892-894. **View Article Google Scholar**

15. Swapna N *et al.* 2017, Knowledge on treatment of Common Minor Ailments among Mothers of under five Children's, Lakshmipuram, Kurnool. *Int J Recent Sci Res.* 8(4), pp. 16615-16617. DOI: http://dx.doi.org/10.24327/ijrsr.2017.0804_0184. **View Article Google Scholar**

16. E. Sree Vani, Katari Kantha, Arumugam Indira, "Knowledge on endotracheal suctioning among staff nurses and nursing students", International Journal of Scientific Research in Science and Technology(IJSRST), Print ISSN : 2395-6011, Online ISSN : 2395-602X, Volume 3, Issue 1, pp.394-397, January-February-2017. **View Article Google Scholar**